

MINISTRY OF HEALTH AND WELLNESS

FACT SHEET FOR DOCTORS RESPONSIBLE FOR THE SENSITISATION OF PUBLIC COMING FOR VACCINATION

FACT SHEET - ASTRAZENECA/COVISHIELD VACCINE

Name	
Surname	
Date	

Persons for whom vaccine is recommended (AstraZeneca / COVISHIELD Vaccine)

	Tick as appropriate (√)
Persons suffering from obesity	
Persons with particular comorbidities :	
o Diabetics	
o Persons with cardiovascular pathologies	
o Persons with respiratory pathologies	
o Persons with renal pathologies, including haemodialysis	
o Persons suffering from cancer	
o Persons with auto immune pathologies	
o Immuno-depressed persons, persons following corticoid treatment and immune-suppressors, persons undergoing chemotherapy	
Persons who have been tested positive for COVID19	

Persons for whom vaccination is not recommended

Persons with history of severe allergy	
Persons with history of allergy following vaccination	
Persons who have had an anaphylactic reaction following first injection of AstraZeneca/COVISHIELD Vaccine	
Vaccination (any vaccination) done 14 days prior to AstraZeneca/COVISHIELD vaccination	
Pregnant women or women planning for pregnancy in the coming month	
Persons aged below 18 (until further notice)	

Persons with whom caution should be exercised

Feverish persons (> 38, 5 degree Celsius) or persons who are subject to an acute infection at the moment of vaccination : these persons shall be recalled after 15 days	
Persons undergoing treatment with an antiplatelet agent shall only be vaccinated through the deltoid muscle. The injection point shall be kept under compress for 5 minutes (should be strictly timed) so as to reduce risks of hematoma at the point of injection	
Vaccination can be done on persons taking anticoagulants under certain particular circumstances, to be verified with a doctor	